

## Lack of Phylogeographic Structure in Nigerian Village Chickens Revealed by Mitochondrial DNA D-loop Sequence Analysis

A.O. Adebambo<sup>1</sup>, V.A. Mobegi<sup>2</sup>, J.M. Mwacharo<sup>2,4</sup>, B.M. Oladejo<sup>1</sup>, R.A. Adewale<sup>1</sup>, L.O. Ilori<sup>1</sup>,  
B.O. Mekanjuola<sup>1</sup>, O. Afolayan<sup>1</sup>, G. Bjørnstad<sup>2,3</sup>, H. Jianlin<sup>2</sup> and O. Hanotte<sup>2,4</sup>

<sup>1</sup>Department of Animal Breeding and Genetics, University of Agriculture,  
Abeokuta, P.M.B. 2240, Abeokuta, Nigeria

<sup>2</sup>International Livestock Research Institute (ILRI), P.O. Box 30709, Nairobi 00100, Kenya

<sup>3</sup>Norwegian School of Veterinary Science, Department of Basic Sciences and Aquatic Medicine,  
P.O. Box 8146 Dep., N-0033 Oslo, Norway

<sup>4</sup>School of Biology, University Park, University of Nottingham, Nottingham NG7 2RD, United Kingdom

**Abstract:** Genetic diversity studies that utilize phenotypic and genetic information are informative when formulating breeding and conservation plans. The present study utilizes sequences of mitochondrial DNA (mtDNA) D-loop region of 232 village chickens from Southern and Northern Nigeria to determine the origin and diversity of Nigerian local chickens. Thirty-six polymorphic sites which generate 35 haplotypes are identified. Phylogenetic analyses group Nigerian local chickens to a single clade and 97.8% of the total maternal variation occurs within populations. Reference sequences representing the major chicken mtDNA lineages from Asia indicate the Indian subcontinent to be the likely main center of origin of Nigerian village chicken. Lack of phylogeographic structure among Nigerian village chickens suggest extensive genetic intermixing within the country.

**Key words:** mtDNA, D-loop, genetic relationship, indigenous village chicken, Nigeria

### INTRODUCTION

Genetic evidence confirms that chickens are derived from multiple maternal origins in Asia (Liu *et al.*, 2006). Chickens are the most widely distributed of all livestock species in Nigeria with a population of 166 million birds (FAOSTAT, 2007). Chicken play very significant socio-cultural and economic roles in most African societies. Quantifying the structure of genetic diversity in different African chicken populations is of significance in optimizing conservation and utilization strategies. The description of Nigerian local chickens is based on phenotypic traits (Nwosu *et al.*, 1985; Adebambo *et al.*, 1999). Such information if complemented with findings obtained using molecular markers could be useful in formulating long term inference/plans for genetic improvement programs. The D-loop region of mitochondrial DNA (mtDNA) has been widely used to assess the diversity and phylogeographic structure of various chicken populations (Niu *et al.*, 2002; Mobegi *et al.*, 2006; Muchadeyi *et al.*, 2008; Razafindraibe *et al.*, 2008). So far, however, no such data has been reported on Nigerian local chicken. In the present study, the hypervariable segment 1 (HV1) region of the D-loop was sequenced to investigate the phylogeographic structure, diversity and origin of Nigerian local chicken.

### MATERIALS AND METHODS

Genomic DNA was extracted from 232 local unrelated chickens from four geographic regions in Nigeria; 58

samples from the Northwest region (NW), 46 from the Northeast region (NE), 96 from the Southwest region (SW) and 32 from the Southeast region (SE). Five hundred and ninety two base pairs of the mtDNA D-loop region were amplified (Mobegi *et al.*, 2005) using L16750 (5'-AGGACTACGGCTTGAAAAGC-3', accession number NC\_001323, Desjardins and Morais 1990) as the forward primer and H547 (5'-ATGTGCCTGACCGAGGAACCAG-3', accession number AB098668, Komiyama *et al.*, 2003) as the reverse primer. Direct sequencing of PCR products was done with the BigDye® Terminator Cycle Sequencing Kit version 3.1 (Applied Biosystems, USA) using two internal primers CR-for (5'-TCTATATTCCACATTCTC-3') and CR-rev (5'-GCGAGCATAACCAAATGG-3'). PCR products were sequenced on ABI 3730 XL capillary sequencer (Applied Biosystems, USA).

A 397 bp long fragment, including the HV1 region was subsequently used for analysis. Following 1000 bootstrap replicates, a Neighbour-Joining (NJ) tree was generated using MEGA 3.0 software (Kumar *et al.*, 2004). Two D-loop sequences of wild junglefowl; *Gallus gallus gallus* and *G. g. bankiva* (GenBank accession numbers AB007720 and AB007718, respectively), were included as out-groups. Seven reference sequences that correspond to the different major clades found in Asian domestic chickens and spanning the geographic range of wild red jungle fowls (Bjørnstad *et al.*, in preparation) were also included in the analysis. The

sequences used were those of the most common haplotype observed in the respective clades. A Median Joining (MJ) network constructed using the program NETWORK 4.1.0.8 (Bandelt *et al.*, 1999) was also generated. Genetic differentiation based on the Analysis of Molecular Variance (AMOVA) and population expansion inferred from mismatch distributions, Tajima's D (Tajima, 1989) and Fu's  $F_s$  (Fu, 1997), were calculated using Arlequin version 3.0 (Excoffier *et al.*, 2005).

## RESULTS

The 232 sequences from local Nigerian chicken populations generated 35 haplotypes from 36 polymorphic sites. All haplotypes have been deposited with the Genbank under accession numbers FJ851656-FJ851686 and GU951751-GU951758 and are shown in Fig. 1. Phylogenetic analysis (Fig. 2) grouped Nigerian local chickens into a single clade (Clade IV) out of the seven clades (Clade I, II, IIIa, IIIb, IIIc, IIId and IV) identified in Asian domestic chicken (Bjørnstad *et al.*, in preparation).




Fig. 1: Nucleotide polymorphisms observed in D-loop HV1 domain of 232 chicken sequences and their frequencies (N). Vertically oriented numbers indicate the site position and the sequences shown are only the variable sites. Dots (.) indicate identity with the reference sequence (GenBank accession number AB098668) (Komiya *et al.*, 2003). The abbreviations denoting the haplotype names are as follows: NE (North East), NW (North West), (NE), SE (South East) and SW (South West)





Fig. 2: Neighbour-joining tree reconstructed using MEGA 3.1 software from 35 haplotypes identified in 232 sequences of Nigerian chicken. Two haplotypes of the genus *Gallus* retrieved from GenBank; *Gallus gallus gallus* (GenBank accession number AB007720) and *Gallus gallus bankiva* (GenBank accession number AB007718) and seven haplotypes representing the major chicken mitochondrial DNA clades (Clade I, II, IIIa, IIIb, IIIc, IIId and IV) are also included in the tree for reference purposes. The numbers at the nodes represent the percentage bootstrap values for interior branches after 1000 replications

Analysis of molecular variance shows that 97.8% of the maternal genetic variation occurs within populations; the remaining variation is found among populations. Figure 3 shows the MJ networks of the sampled Nigerian local chicken populations. As in the phylogenetic tree, the 35 haplotypes clustered together in Clade IV. Figure 4



is chicken based on the polymorphic  
proportional to the frequency of the  
refer to the positions of nucleotide  
(r AB098668)

theory of population expansion among  
 ens. Positive values shown by SSD  
 Raggedness index  $r$  also indicate  
 tion of new and low-frequency  
 ation variation typical in expanding

rent study indicate that mtDNA (HV1)  
 highly variable in Nigerian local  
 a total of 35 haplotypes. All the  
 uences observed in Nigerian village  
 from a single sequence (NE84)  
 IV, which likely originates from the  
 t (Bjørnstad *et al.*, in preparation).  
 06) also observed that majority of  
 in West African village chicken  
 in clade IV, while Muchadeyi *et al.*  
 indraibe *et al.* (2008) observed two  
 es amongst both Zimbabwean and

ary of population expansion among  
ens. Positive values shown by SSD  
Raggedness index  $r$  also indicate  
ution of new and low-frequency  
ation variation typical in expanding

present study indicate that mtDNA (HV1) is highly variable in Nigerian local chickens, with a total of 35 haplotypes. All the haplotypes observed in Nigerian village chickens are derived from a single sequence (NE84) in clade IV, which likely originates from the West African chicken (Bjørnstad *et al.*, in preparation). Muchadeyi (2006) also observed that majority of haplotypes in West African village chickens are in clade IV, while Muchadeyi *et al.* (2006) and Muchadeyi *et al.* (2008) observed two haplotypes amongst both Zimbabwean and




Fig. 4: Geographic distribution of the 35 haplotypes observed in the D-loop HV1 segment of the 232 Nigerian chicken sequences




Fig. 5: Mismatch distribution of Nigerian indigenous chicken. Observed and model frequencies are indicated

Malagasy chickens, indicating a different history of domestic chicken between the West African and the Southern African regions. Fig. 3 and Fig. 4 indicate no clear geographic sub-structuring in Nigerian chicken population, likely a consequence of the relative recent arrival of domestic chicken in the region and the lack of phylogeographic resolution of the marker used in this study. Alternatively, it could equally be the result of intensive genetic intermixing between populations following human migrations and trading. Haplotype NE84 shows the widest geographic distribution, but with an increasing frequency in the Southwest direction. This possibly infers a North-South dispersal of this haplotype. The D-loop region in Nigerian chicken shows a departure from equilibrium (significant negative values

for Tajima's  $D$  and Fu's  $F_s$ ) suggesting population expansion. These results are consistent with the observed mismatch distribution patterns and non-significant value for Harpending's raggedness index. Therefore, a population expansion of Nigerian local chicken likely followed their arrival in the country. The lack of phylogeographic structure observed among the different populations sampled from different geographic regions suggests population intermixing across long distances within the country.

The ancestors of the today's Nigerian chicken could have reached the country from the North of the African continent through trans-Saharan trading and/or could have originated from East Africa through migrations along the Sahelian belt. Alternatively, they could have also reached Nigeria from the coast, e.g. following early European exploration. Our data do not allow us to distinguish between these hypotheses but the relative simple network expansion observed, characterized by an expansion from a single ancestral haplotype (Fig. 3) suggests a single geographic origin. Whatever the case, it appears that the Asian maternal ancestor of Nigerian local chicken would have originated initially from the Indian subcontinent as all haplotypes identified in local Nigerian chicken belong to clade IV which most likely originates from the Indian subcontinent (Bjørnstad *et al.*, in preparation).

**Conclusion:** Overall, Nigerian local chicken populations would have likely been introduced first into the Northern parts of the country. Thereafter the chicken populations

dispersed towards the southernmost region following human movements and commercial activities. The lack of substructure in the chicken populations is evidence of a single maternal origin and extensive genetic intermixing in the past and present times.

## ACKNOWLEDGEMENTS

The authors would like to acknowledge the assistance provided by the members of staff in the Biotechnology theme (International Livestock Research Institute, Nairobi, Kenya); the Biotechnology Center and Department of Animal Breeding and Genetics, University of Agriculture, Abeokuta, Nigeria.

## REFERENCES

- Adebambo, O.A., C.O.N. Ikeobi, M.O. Ozoje and J.A. Adenowo, 1999. Color variations and performance characteristics of the indigenous chicken of South Western Nigeria. *Nig. J. Anim. Prod.*, 26: 15-22.
- Bandelt, H.J., P. Forster and A. Rohl, 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.*, 16: 37-38.
- Bjørnstad *et al.* (unpublished data). Multiple maternal origins for the domestic chicken.
- Desjardins, P. and R. Morais, 1990. Sequence and gene organization of the chicken mitochondrial genome. A novel gene order in higher vertebrates. *J. Mol. Biol.*, 212: 599-634.
- Excoffier, L., G. Laval and S. Schneider, 2005. ARLEQUIN ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics*. Online, 1: 47-50.
- FAOSTAT, 2007. Food and Agricultural Organization statistical databases. CDROM.
- Fu, Y.X., 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics*, 147: 915-925.
- Komiyama, T., K. Ikeo and T. Gojobori, 2003. Where is the origin of the Japanese gamecocks? *Gene*, 317: 195-202.
- Kumar, S., K. Tamura and M. Nei, 2004. MEGA3: Integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform*, 5: 150-163.
- Liu, Y.P., G.S. Wu, Y.G. Yao, Y.W. Miao, G. Luikart, M. Baig, A. Beja-Pereira, Z.L. Ding, M.G. Palanichamy and Y.P. Zhang, 2006. Multiple maternal origins of chickens: out of the Asian jungle. *Mol. Phylogenet. Evol.*, 38: 12-19.
- Mobegi, A.V., 2006. Chicken Diversity Consortium. Mitochondrial DNA D-loop Sequences Reveal the Genetic Diversity of African Chicken. In: Rege, J.E.O., A.M. Nyamu and D. Sendalo (Eds.). Tanzania Society for Animal Production, Dar es Salaam; All Africa Society for Animal Production, Nairobi (Kenya). The role of biotechnology in animal agriculture to address poverty in Africa: opportunities and challenges. Proceedings of the 4th all Africa conference on animal agriculture and the 31st annual meeting of the Tanzania Society for Animal Production. Dar es Salaam (Tanzania): TSAP, pp: 293-298.
- Muchadeyi, F.C., H. Eding, H. Simianer, C.B.A. Wollny, E. Groeneveld and S. Weigend, 2008. Mitochondrial DNA D-loop sequences suggest a Southeast Asian and Indian origin of Zimbabwean village chickens. *Anim. Genet.*, 39: 615-622.
- Niu, D., Y. Fu, J. Luo, H. Ruan, X.P. Yu, G. Chen and Y.P. Zhang, 2002. The origin and genetic diversity of Chinese native chicken breeds. *Biochem. Genet.*, 40: 163-174.
- Nwosu, C.C., F.A. Gowen, F.C. Obioha, I.A. Akpan and G.I. Onuora, 1985. Biometrical study of the conformation of the native chicken. *Nig. J. Anim. Prod.*, 12: 141-146.
- Razafindraibe, H., V.A. Mobegi, S.C. Ommeh, Rakotondravao, G. Bjørnstad, O. Hanotte and H. Jianlin, 2008. Mitochondrial DNA origin of indigenous Malagasy chicken: implication for a functional polymorphism at the Mx gene. *Ann. N.Y. Acad. Sci.*, 1149: 77-79.
- Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123: 585-595.